



## AWARENESS AND UTILIZATION PATTERN OF *JATROPHA CURCAS*(L) IN YEWA SOUTH LOCAL GOVERNMENT AREA, OGUN STATE, NIGERIA.

Adedokun, M. O., Adegbenjo, A. E., Matanmi A. S., and Soaga J. A.

Department of Forestry and Wildlife Management,  
Federal University of Agriculture, Abeokuta.  
Correspondence E-mail: [ayanyemiadegbenjo@gmail.com](mailto:ayanyemiadegbenjo@gmail.com)

Adedokun, M. O., Adegbenjo, A. E., Matanmi A. S. and Soaga J. A. (2018): Awareness and Utilization Pattern of *Jatropha Curcas*(L) in Yewa South Local Government Area, Ogun State, Nigeria. *Nigerian Journal of Forestry*, 48 (2) 46 - 50

### Abstract

This study investigated the utilization pattern of *Jatropha curcas* plant and phytochemical properties of *Jatropha curcas*(L) from Yewa South Local Government Area, Ogun State, Nigeria. A purposive sampling technique was used in selecting 120 respondents and data were collected from them through questionnaires. Data analyses were carried out and result presented using descriptive statistics. Information were sought on level of awareness about *Jatropha curcas* and level of utilization of its derivatives. Findings revealed that the respondents are vast in the use of *Jatropha curcas* for traditional medicine. The study also showed that *Jatropha curcas* has an average oil yield of 32% hence, it is efficient for use as bio-fuel. The highest phytochemical detected was alkaloids (8.1%). Biofuel has been regarded as potential alternative for fuel for partial substitution of bio-fuel. *Jatropha curcas*, which is one of the sources of bio-diesel, is readily available in tropical and sub-tropical areas. Awareness of *Jatropha curcas* plant usefulness but not that of its product is very high in Yewa south local government area. Deliberate promotion of the plant products and the establishment of *Jatropha* System for sustainable production among the populace is therefore recommended.

**Keywords-** *Jatropha curcas*, Biofuel, Phytochemical, Utilization, Awareness

### Introduction

The *Jatropha curcas* L. belongs to the family Euphorbiaceae and the genus *Jatropha*. It is considered to have originated from Central America, most probably Mexico, where it is found extensively in different regions of the country (Enzo, 2007). *Jatropha* plants that occurred in various climate regions and sometimes from the same climate zone shows morphological differences particularly with regards to their shape, size of seeds, protein and lipid content (Jorge *et al.*, 2010). *J. curcas* is a deciduous shrub that grows up to a height of 3.5 meters and with a production life of 50 years (Azza and Ferial, 2010) The genus *Jatropha* includes a large number of species (170) that are distributed over the world in habitats ranging from the tropical to the temperate zone (Enzo, 2007).

It is fast growing plant and produces seeds after approximately 1-3 years, (Jorge *et al.*, 2010). depending on rainfall conditions and how the plant is propagated. *Jatropha* grows on well-drained soil with good aeration and is well adapted to marginal soil with low nutrient content (Azza and Ferial, 2010). It occurs mainly at lower altitudes (0-500m) in areas with average annual temperatures well above 20°C but can grow at higher altitudes and tolerates slight frost (Augustus *et al.*, 2002).

The Arabs have been using this plant for medicinal purpose (Enzo, 2007). Today it is found in almost all the tropical and sub-tropical regions of the world (Jorge *et al.*, 2010). There are more than 200 different names

for its great significance to man and the various possibilities of its use The botanist Carl Von Linne first classified the plants in 1753, gave it the botanical name "*Jatropha curcas*" from the Greek word "Jatros" meaning a "Doctor" and "trophe" meaning "nutrition" (Augustus *et al.*, 2002). Even Linne had realized the potential of this plant for medicinal purposes (Gubitz *et al.*, 1999). The plant is regarded as a shrubs/small tree with height generally ranging from 3-5 meter (m). It has been estimated that the life of the plant is up to 50 years. Different varieties of the plant can be found which are generally Cape Verde, Nicaragua, Ife- Nigeria, non-toxic Mexico (The Biomass Project, 2000).

*Jatropha* is a valuable multipurpose crop that alleviates soil degradation, desertification and deforestation which can be used for bio-energy to replace petrodiesel, soap production and climate amelioration protection (CJP 2007). *Jatropha* It is a small tree with smooth gray bark, which exudes whitish colored watery latex cut and its stain is difficult to remove (Enzo, 2007).

Normally it grows between (3-5m) in height, but can attain a height of up to (8-10m) with diameter up to 20cm under favorable conditions (Henning, 2000).

The recent interest as alternative sources of energy placed *J. curcas* in the fore front of discussions on bio-fuel production. Apart from biofuel, it also produces wood (though of poor quality) for cooking, animal feeds and is used for rehabilitation of marginal lands (Heller, 1996). These characteristics coupled with the hoax towards energy independence placed *J. curcas* in a

1996). These characteristics coupled with the hoax towards energy independence placed *J. curcas* in a strategic position for developing countries, especially Nigeria, where the demand for fossil fuel is not marched by a corresponding increase in the supply infrastructure. Therefore, this study aimed to access the utilization pattern and phytochemical properties of *J. curcas* seed and the oil.

**Materials and Method**

The study was carried out in Yewa South Local Government area of Ogun State, Nigeria. It lies on (6°48'N 2°57'E) and has land mass of 163,720 square hectares and a population of 168,850 according to 2006 census (Wikipedia, 2008). A purposive sampling method was adopted in selecting Ilaro, Idogo, Okeodan, Ajilete and Ilobi areas of Yewa south local Government area of Ogun state, Nigeria, due to the abundance of *J. curcas* in the area. A well-structured questionnaire was administered to 24 respondents in each the selected areas and *J. curcas* seeds were also collected for oil extraction and phytochemical analyses. The matured seeds of *J. curcas* were collected from free areas within Ilaro in Yewa south

local Government area, Ogun State, Nigeria. The seed was de-hulled manually and separated by winnowing, it was grinded into smaller pieces using a grinding stone and dried under room temperature. The oil was extracted at the Nigerian Institute of Science Laboratory Technology (NISLT, Ibadan) using a soxihet extractor and normal-hexane as the reagent or solvent, the soxihet extractor has three main parts which is the thimble, round bottom flask and the condenser. The thimble was loaded with the sample (grinded *Jatropha curcas* seed), the rounded bottom flask was loaded with the reagent/solvent which is normal-hexane and heated (70°C) which aids in syphoning oil content from the sample and the condenser has an enlarged uptake tube to improve condensing capacity after about 6 hours the round bottom flask contain both the solvent and oil extracts forming a solution. Then, it was taking to the oven where it was heated at the boiling point of the solvent used (normal-hexane) 69°C leaving concentrated oil extract, the phytochemical properties analysis of the oil was then carried out. The total seeds collected from the area weighed 372g while the quantity of oil extracted was 120g

$$\text{Oil yield (\%)} = \frac{\text{Quantity of oil extract}}{\text{Quantity of seed}} \times 100$$

$$\text{Oil yield (\%)} = \frac{120g}{372g} \times 100 = 32\%$$

**Result and Discussion**

**Table 1- Socio-Economic Characteristics of the Respondents**

<b>Variables</b>	<b>Frequency</b>	<b>Percentage (%)</b>
<b>Age</b>		
Below 26	5	4.2
26-35	30	25
36-45	33	27.5
Above 45	52	43.3
<b>Total</b>	<b>120</b>	<b>100</b>
<b>Gender</b>		
Male	68	56.7
Female	52	43.3
<b>Total</b>	<b>120</b>	<b>100</b>
<b>Marital status</b>		
Single	23	19.2
Married	63	52.5
Others	34	28.3
<b>Total</b>	<b>120</b>	<b>100</b>
<b>Occupation</b>		
Farming	43	35.8
Civil servant	35	29.2
Trading	31	25.8
Others	11	9.2
<b>Total</b>	<b>120</b>	<b>100</b>

Table 1 showed the socio-economic characteristics of the respondents. From the result, 43.3% of the respondent were above 45 years, 27.5% were between 26 - 35 years, 25% were between 36 - 45 years and 4.2% were below 26 years. Since majority of the

respondent are adults, this could positively affect their vastness in the knowledge of *J.curcas* plant and its derivatives. The result also revealed that 56.7% were male while 43.3% were female. Marital status indicated that 52.5% were married, 19.2% are singles and 28.3% are others (Widowed).

**Table 2: Percentage Distribution on Awareness and Utilization of *J.curcas* Plant and its Derivatives**

Variables	Frequency	Percentage (%)
<b>Do you know <i>Jatropha curcas</i> plant?</b>		
Yes	114	95
No	6	5
Total		
<b>Do you use any part of the plant?</b>		
Yes	86	71.7
No	34	28.3
Total	120	100
<b>Which part of the plant do you use</b>		
Leaves	21	24.4
Root	16	18.6
Seeds	9	10.5
Stem	40	46.5
Total	86	100
<b>Leaves</b>		
Gonorrhoea	6	28.6
Staphylococcus	6	28.6
Bathing infant to avoid sickness	9	42.9
<b>Root</b>		
Malaria/fever	16	100.0
<b>Seed</b>		
Inducing child bearing	9	100.0
<b>Stem</b>		
Chewing stick		
<b>What do you use it for</b>		
Fuel	1	1.2
Condiment	1	1.2
Medicine/curing diseases	84	97.6
Total		
<b>Do you know if the seed has oil?</b>		
Yes	8	88.9
No	1	11.1
Total	9	100
<b>Have you extracted oil from the seed?</b>		
Yes	3	33.7
No	6	66.3
Total	9	100
<b>Method used in extracting the oil</b>		
Local	2	33.7
Modern	1	66.3
Total	3	100
<b>What do you use the oil extracted for?</b>		
Fuel	1	33.7
Medicinal purpose	2	66.3
Total	3	100

In Table 2, majority of the respondents (95%) in the study areas had knowledge of the species utilization. The level of its utilization among the respondents is also shown in Table 2. It can be deduced that 71.7%

uses parts of *Jatropha curcas* plant, only 23.3% do not use the plant parts. From the sampled-respondents, it was revealed that 24.4% uses the leaf for curing of diseases like gonorrhoea, staphylococcus and it is also

used in bathing infants so as to avoid sickness, 18.6% uses the root for curing malaria and fever, 10.5% uses the seed for inducing child birth in women by curing vaginal infection and it is also used as fuel in company with its abortifacient potential, and majority uses the stem (46.5%) mainly for chewing stick. It also reveals that 1.2% use it as a fuel source, 1.2% uses it as condiments and majority (87.6%) of the respondents utilizes it for medicinal purposes and treatment of ailments.

The result further showed that a larger population (88.9%) of the respondents that utilizes the seed of *Jatropha curcas* were aware of its soil content, while

11.1% were not. Only 33.3% of respondents know how the oil is extracted from its seeds and 66.7% has never extracted the oil. However, local method by boiling with water for hours after which it is further subjected to purification methods were adopted by most of the respondents in the extraction of oil from the seed. Essentially, respondents that extracted the oil showed that extracted oil from the seeds is used for medicinal purposes (66.7%) and 33.7% of the respondents claimed its use for fuel. The research discovered that the oil yield of *Jatropha curcas* was 32% which is fairly average compared to 20-40% reported by Augustus *et. al.*, 2002.

**Table 3- Phytochemical Properties of *J. curcas* Seed Oil**

Phytochemicals	Values from seed oil
Flavanoid	1.3%
Alkaloids	8.1%
Oxalate	0.181mg/g
Phytic acid	1.01%
Tannin	0.17%
Saponin	0.01%
Phenol	0.75%

Low concentration of flavonoids (1.3%), alkaloids (8.1%) which is very high compared to 1.01% seed oil recorded by (Ijei and Omodairo, 2006), and tannin concentration of (0.17%) were observed in *J. curcas* seed oil (Table 3), tannin has anti-inflammatory effect which help control all indication of gastritis, esophagitis, enteritis and irritating bowel disorder (Dhannananda, 2001), tannins are also responsible for antidiarrheal activity (Enzo, 2007). These values are low and may be considered as of no nutritional significance. The phytic acid content of *Jatropha curcas* seed was 1.01%. This value is close compared with 1.5% for soybean reported by Gubitzet *al.*, (1999). Saponins concentration in *J. curcas* was lower than other anti-nutritional factors under study, saponin are used as dietary supplements, expectorant and anti-inflammatory agent (Enzo, 2007). Phenolic compound in *Jatropha curcas* seed oil have antiseptic properties which could explain the use of this oil for the treatment of various diseases (Azza and Ferial, 2010). The antimicrobial activities of phytochemicals are further evidences by their active role in plant disease resistance (Ijeh&Omodairo, 2006) and antioxidant activity. *Jatropha curcas* plant is widely used as an application to bruises – recast and wounds in herbal medicine in western Nigeria (Gubitzet *al.*, 1999). The exudates of *Jatropha curcas* effectively arrested bleeding from fresh wounds, inhibited microbial growth of known wound contaminants and accelerates wound healing process (Gubitzet. *al.*, 1999).

**Conclusion and Recommendation**

Potential use of *J. curcas* as an energy crops, medicinal plant, agro-forestry crops, soil conservation measures and industrial application are the attractive factors to cultivate the plant in the unused and barren land.

However, awareness of the utilization of its products and the opportunities it offers in terms of income and employment generation is very low. Planting of *J. curcas* by the respondents is not as a result of their awareness about biodiesel, but because it is an age long activity for boundaries marking. Multiple benefits of *J. curcas* plants and oil expelled from its seeds are not only useful in saving the environmental pollution but supports for employment generation and entrepreneurship developments. The results of this investigation suggest that *Jatropha curcas* seed oils have great potentials in the prevention and treatment of many ailments and also shows presence of some toxic substance hence cannot be digested by human hence it should be subjected to further purification if it is needed to be used for cooking.

**References**

Augustus G.D.P.S., Jayabalan M., and Seiler G.J. (2002): Evaluation and bio-induction of energy components of *Jatropha curcas*. Biomass and Bioenergy, Vol.23, No. 3, (September 2002), pp. 161-164, ISSN: 0961-9534

Azza, A.A. and Ferial, M.A. (2010). Nutritional Quality of *Jatropha curcas* seeds and effect of some physical and chemical treatments on their anti-nutritional factors. African Journal of Food Science. 4: 93-103. Centre for *Jatropha* Promotion and Biodiesel, (2007). Handbook B-132, Sainik Basti, Churu-331001, Rajasthan, India, pp.133

Dharmananda, S. (2001). Gallnuts and the use of tannin in Chinese medicine. Institute of traditional medicine, Portland, Oregon, pp 50-57

- Enzo A.P., (2007). Traditional plants and herbal remedies used in the treatment of diarrhea, Diseases: Mode of action, quality, efficacy and safety considerations. Modern phytochemical turning medicinal plant to drugs. WILEY.VCH Verlag GmbH and Co.KGaA, Weinheim, pp. 248-260
- Gubitz G.M., Mittelbach M., and Trabi M. (1999) Exploitation of tropical oil seed plant of *Jatropha curcas*L., Bioresources Technology; 67, 73-82
- Heller J. (1996): Physics nut- *Jatropha curcas*L. Promoting the conservation and use of underutilized and neglected crops (1). Institute of Plant Genetics and Crop Plant Research, Gatersleben International Plant Genetics Resource Institute, Rome. 66 pp.
- Henning Richard K. The *Jatropha* Booklet, published in the *Jatropha* website, [www.jatrophpha.de/documents/jcl-booklet.pdf](http://www.jatrophpha.de/documents/jcl-booklet.pdf)(2000)
- Ijeh, I.I., and Omodairo, O.J. (2006). Anti-microbial effect of aqueous and ethanoic fractions of some local spices. *Ocimumgrassitum* and *Xylophia aethiopica*. Recent Progress in medicinal plant. 13: 455-460.
- Jorges M.H., Harinder M., Gorge F., and Klaus B. (2010). Agroclimate conditions, Chemical and Nutritional Characterization of different provinces of *Jatropha curcas*L from Mexico 6 (14): 1690-1696.
- Reddy, N.R., and Pierson, M.D. (1994). Reduction in anti-nutritional and toxic components in plant foods by fermentation. Food Research International 27, 281-290
- The Biomass Project, (2000). "Curcas Oil Methyl Ester Nigaragua <http://www.ibw.com.ni/biomasa/mat.htm> (2008/01/09)
- Wikipedia, 2008. Oyo State. Available on <http://en.wikipedia.org/wiki/Oyostate>. Accessed 10/10/2015